

Race, Treatment Preferences, and Hospice Enrollment: Eligibility Criteria May Exclude Patients With the Greatest Needs for Care

Jessica Fishman, PhD^{1,2,3}, Peter O'Dwyer, MD³, Hien L. Lu, BA³, Hope Henderson, PA⁴, David A. Asch, MD, MBA^{3,4,5}, and David J. Casarett, MD, MA^{3,4,5}

BACKGROUND: The requirement that patients give up curative treatment makes hospice enrollment unappealing for some patients and may particularly limit use among African-American patients. The current study was conducted to determine whether African-American patients with cancer are more likely than white patients to have preferences for cancer treatment that exclude them from hospice and whether they are less likely to want specific hospice services. **METHODS:** Two hundred eighty-three patients who were receiving treatment for cancer at 6 oncology clinics within the University of Pennsylvania Cancer Network completed conjoint interviews measuring their perceived need for 5 hospice services and their preferences for continuing cancer treatment. Patients were followed for 6 months or until death. **RESULTS:** African-American patients had stronger preferences for continuing their cancer treatments on a 7-point scale even after adjusting for age, sex, finances, education, Eastern Cooperative Oncology Group performance status, quality of life, and physical and psychologic symptom burden (adjusted mean score, 4.75 vs 3.96; β coefficient, 0.82; 95% confidence interval, 0.22-1.41 [$P=.007$]). African-American patients also had greater perceived needs for hospice services after adjusting for these characteristics (adjusted mean score, 2.31 vs 1.83; β coefficient, 0.51; 95% confidence interval, 0.11-0.92 [$P=.01$]). However, this effect disappeared after adjusting for household finances. **CONCLUSIONS:** Hospice eligibility criteria may exclude African-American patients disproportionately despite greater perceived needs for hospice services in this population. The mechanisms driving this health disparity likely include both cultural differences and economic characteristics, and consideration should be given to redesigning hospice eligibility criteria. **Cancer 2009;115:689-97.** © 2008 American Cancer Society.

KEY WORDS: treatment preferences, conjoint analysis, utilities, interview, Medicare Hospice Benefit, services, palliative care, African American, race, economic, finances.

Patients with cancer have substantial needs for care throughout the course of their illness, from diagnosis through the end of life.¹⁻⁸ Hospice offers a comprehensive program of services designed to meet these

Corresponding author: David J. Casarett, MD, MA, University of Pennsylvania, 3615 Chestnut Street, Philadelphia, PA 19104; Fax: (215) 573-8684; casarett@mail.med.upenn.edu

¹Annenberg School for Communication, University of Pennsylvania, Philadelphia, Pennsylvania; ²Center for Clinical Epidemiology and Biostatistics, University of Pennsylvania, Philadelphia, Pennsylvania; ³University of Pennsylvania School of Medicine, Philadelphia, Pennsylvania; ⁴Veterans Affairs Center for Health Equity Research and Promotion, Philadelphia, Pennsylvania; ⁵Leonard Davis Institute of Health Economics, University of Pennsylvania, Philadelphia, Pennsylvania

Received: June 16, 2008; **Revised:** August 5, 2008; **Accepted:** September 2, 2008

Published online: December 22, 2008, © 2008 American Cancer Society

DOI: 10.1002/cncr.24046, www.interscience.wiley.com

needs, including a visiting nurse, respite care, a chaplain, a home health aide, and a counselor. Approximately 500,000 patients with cancer enroll in hospice every year, and evidence indicates that hospice enrollment is associated with improved patient and family outcomes and higher satisfaction with care.⁹⁻¹⁴

African-American patients are less likely than white patients to use hospice.¹⁵⁻¹⁷ Although these disparities are well described, it is not known why they exist. It is possible that African Americans delay or avoid hospice enrollment because of the requirement that they forgo curative treatment.¹⁸ This criterion forces patients to make a 'terrible choice' between continued treatment and hospice services,¹⁹⁻²¹ a choice widely believed to obstruct timely hospice use.²²⁻²⁵ The criterion has been part of the Medicare Hospice Benefit since its creation 25 years ago; and, because nearly all hospice are Medicare-certified, they follow Medicare eligibility criteria for all patients regardless of age or insurance. This barrier may be particularly significant for African Americans, who are more likely than whites to want various life-prolonging interventions.²⁶⁻²⁹ However, it is not known whether African-American patients also have more aggressive cancer treatment preferences or whether such preferences help explain the disparity in hospice use.

An alternative explanation for the under-use of hospice among African Americans is that they may be less likely to want the services that hospice provides. To our knowledge, no studies have evaluated this possibility. Although some studies suggest that African Americans have less favorable attitudes toward hospice,³⁰⁻³⁴ their perceived needs for hospice services have not been described.

It is important to determine why these disparities in hospice enrollment exist.^{35,36} If they are the result of aggressive treatment preferences among African Americans, then lower rates of hospice use, arguably, are unfair and represent a real disparity in care that could be eliminated by redesigning hospice eligibility criteria. Conversely, if African Americans are less likely to want hospice services, then changes to the benefit are not necessary, and, instead, modifications to the services offered may be warranted. Therefore, the objective of this study was to define and compare preferences for cancer treatment and perceived needs for hospice services among African-American patients and white patients.

MATERIALS AND METHODS

Setting and Sample

We conducted this study in a population of patients with cancer, because cancer is the most common hospice admitting diagnosis (approximately 50%).³⁷ Patients with cancer were recruited over 18 months from 6 oncology clinics within the University of Pennsylvania Comprehensive Cancer Center Network. Patients were eligible if they had clinical or radiologic evidence of active cancer, were receiving chemotherapy or radiation therapy, and had a life expectancy of ≤ 6 months if they were to discontinue cancer-directed treatment, according to their oncologist. These patients met the prognostic eligibility criterion for hospice, which requires that patients have a prognosis of ≤ 6 months if their disease runs its usual course (eg, without treatment).¹⁸ Nursing staff at each clinic assisted a rotating team of 3 interviewers to identify and approach eligible patients. This study was approved by the institutional review boards of participating sites.

Data Collection

After providing informed consent, patients completed separate structured interviews using Tablet personal computers. First, they provided demographic data, including self-described race, age, marital status, and household finances (eg, money available at the end of the month).³⁸ In addition, patients completed the Global Distress Index (GDI) of the Memorial Symptom Assessment Scale,³⁹ the Functional Assessment of Cancer Therapy-General (FACT-G),⁴⁰ the Medical Outcomes Survey Social Support Scale,⁴¹ a single-item global rating of health,⁴² the Eastern Cooperative Oncology Group performance status scale (ECOG-PS),^{43,44} and assessments of Activities of Daily Living (ADLs) and Instrumental Activities of Daily Living (IADLs).^{45,46}

Patients' perceived needs for hospice services were assessed using a conjoint interview, which refers to a process by which respondents are asked to make choices among goods or services by considering several attributes jointly.⁴⁷ One advantage of this approach over direct ratings is that the value of each service is determined while considering other services, revealing preferences in a more naturalized setting rather than eliciting them 1 by 1. Second, whereas direct rating tasks often suffer from ceiling

Nurse: A nurse monitors and explains the patient's condition and could help reduce any symptoms the patient may be having, such as pain or nausea. A nurse helps manage patient medications, and could answer questions that patients and families have.

Chaplain: A chaplain provides spiritual and emotional support and can discuss concerns about the future.

Home health aide: A home health aide provides help with personal needs, such as bathing and dressing, housework, cleaning, and laundry.

Counselor: A counselor is available to talk about concerns about the patient's illness and can identify depression, anxiety or other conditions the physician should know about.

Respite Care: Respite care offers a place that a patient can go to stay for a few days to give family members or other caregivers a rest.

FIGURE 1. Descriptions of hospice services.


effects, in conjoint tasks, respondents must choose those services that are most important.

For these interviews, we used Sawtooth Software's Adaptive Conjoint Analysis (ACA) package (ACA for Windows, version 4.0; Sawtooth Software, Sequim, Wash). ACA is a hybrid⁴⁸⁻⁵⁰ method of conjoint analysis that uses an interactive, self-administered computer program to decrease respondent burden while preserving the psychometric advantages of conjoint techniques. Similar to classic conjoint methods, hybrid conjoint methods such as ACA determine the importance that respondents place on various services by examining the choices that they make between pairs of services. In addition, however, hybrid methods also use respondents' direct ratings of the importance of those services, which substantially reduces the number of choices that each individual is required to make. Although they were developed for use in marketing,⁴⁷ conjoint techniques and hybrid modifications have been used increasingly to study health-related preferences.⁵¹⁻⁵⁸ Their validity, reliability, and predictive power are well established.^{59,60}

Patients reviewed a brochure describing 5 of the hospice services that are required by Medicare: a visiting nurse, a chaplain, a home health aide, a counselor, and respite care¹⁸ (Fig. 1). These services were selected based on previous studies indicating that they are among the most valuable to patients.⁶¹⁻⁶³ The subsequent interview used abbreviated (4-6 word) descriptions of each service, and patients kept the brochure open for rapid reference. To assess patients' perceived needs for each service, they were asked to consider how much each service would help

Direct ratings of importance

If two home care service plans were acceptable to you in all other ways, how important would this difference be in your enrollment decision?



Pairs questions

If these two home care service plans were exactly the same in every other way, which would you prefer?




FIGURE 2. Examples of the questions used in conjoint interviews to measure patient's perceived needs (direct ratings of importance and comparisons of pairs of services).

them if it were offered at that time (Fig. 2). First, they rated the importance of each service on a scale from 1 ('not at all important') to 7 ('extremely important'). Second, they evaluated pairs of programs presented side-by-side, each containing 2 or 3 services, and rated their preference on a scale from 1 to 9 (1, 'strongly prefer the program on the left'; 5, 'indifferent'; 9, 'strongly prefer the program on the right').

Patients' preferences for cancer treatment were assessed by asking whether they would want to continue receiving their current cancer treatment to achieve various probabilities of surviving for 6 months (nearly 100%, 90%-99%, 50%-89%, 10%-49%, 1%-9%, or nearly 0%). This question was adapted from previous work in seriously ill populations.^{64,65} These probabilities were varied systematically (from lowest to highest and then back to the second lowest) to balance order effects. The lowest probability of 6-month survival was recorded for which a patient would be willing to continue receiving treatment. Responses were used to create a 7-level ordinal variable that described the strength of a patient's treatment preferences (1 level for each of the 6 responses above plus 1 for those patients who would not want to continue treatment even for a nearly 100% chance of surviving for 6 months).

This last category was used to create a dichotomous variable that identified patients who would not want treatment even for a nearly 100% probability of surviving for 6 months and, thus, who would have been eligible for hospice.

Data Analysis

By using the scale described above, preferences for aggressive cancer treatment were compared using a Wilcoxon rank-sum test in bivariate analysis. Adjusted analysis of preferences was performed by using an ordinal logistic regression model. On the basis of respondents' direct ratings and their choices among service options, ACA uses least-squares regression to calculate the utility, or usefulness, of each service for each patient.⁶⁶ These utilities reflect the perceived need that a patient has for each service, and higher utilities correspond to a greater perceived need. We analyzed patients' perceived needs for each of the 5 services individually as well as patients' perceived needs for all 5 services combined (their total utilities). Because utilities were not distributed normally, we used the Wilcoxon rank-sum test to compare perceived needs among patient subgroups and linear regression models with a log transformation and exponentiated coefficients to identify variables that were associated independently with service needs. Variables with a significance value $<.25$ were considered for inclusion in a multivariable linear regression model,⁶⁷ also with a log transformation and exponentiated coefficients.

A sample of 283 patients would be adequate to detect a small difference in the summed utilities of all 5 hospice services between African-American and white patients, assuming at least 20% of the sample would be African American (ability to detect 0.33 standardized difference, $\alpha = .05$; 2-sided). Stata software (Stata for Windows, version 8.0; StataCorp, College Station, Tex) was used for all statistical analysis.

RESULTS

Patient Characteristics

Patients who agreed to participate (300 of 352; 85%) were similar to those who refused with respect to age, ECOG-PS, race, and education (data not shown). How-

ever, women were significantly more likely to participate compared with men (152 of 169 women [92%] vs 148 of 183 men [82%]; Fisher exact test; $P = .02$). Of the 300 patients who consented, this sample was restricted to those patients ($n = 283$; 94%) who self-identified as either African-American ($n = 81$; 29%) or white ($n = 202$; 71%). The characteristics of these patients are described in Table 1.

Patient Race and Cancer Treatment Preferences

African-American patients had stronger preferences for aggressive cancer treatment (using the scale of choices from 1 to 7) compared with white patients (median score, 6 vs 4; Wilcoxon rank-sum test; $P = .006$). That is, they were willing to continue their treatment in return for a smaller likelihood of 6-month survival. In an ordinal logistic regression model, African-American patients had stronger preferences for aggressive cancer treatment (adjusted means, 4.75 vs 3.96; β coefficient, 0.82; 95% confidence interval [95% CI], 0.22-1.41 [$P = .007$]) after adjusting for age, sex, household finances, education, ECOG-PS, quality of life (FACT-G), and physical and psychologic symptom burden (physical and psychologic subscales of the GDI, respectively). Compared with white patients, African-American patients also were less likely to say that they would refuse all cancer treatment even if it were associated with an 'almost 100%' probability of surviving for 6 months (5 of 81 patients [6%] vs 32 of 202 patients [16%]; odds ratio, 2.86; 95% CI, 1.04-9.74; $P = .03$).

Patient Race and Perceived Needs for Hospice Services

Compared with white patients, African-American patients reported greater needs for all hospice services combined (median utilities: 2.26 vs 1.79; rank-sum test; $P = .005$). African-American patients assigned higher utilities to all hospice services except a home health aide (Fig. 3).

In a multivariate regression model, African-American race remained associated with greater perceived needs for services after adjusting for age, education, ECOG-PS, dependencies in ADLs, and physical and psychologic symptom burden (GDI physical and psychologic

Table 1. Patient Characteristics

Characteristic	No. of Patients (%)			P
	Overall, N=300	African American, N=81	White, N=202	
Sex				.10
Men	148 (49)	47 (58)	95 (47)	
Age, y				.002
Mean [range]	58 [20-89]	63 [31-89]	57 [20-89]	
>65	84 (28)	33 (41)	49 (24)	
Marital status				.20
Single	45 (15)	19 (23)	23 (11)	
Married	174 (58)	28 (35)	136 (67)	
Divorced/separated	45 (15)	19 (23)	23 (11)	
Widowed	22 (7)	9 (11)	13 (6)	
Living w/ partner	14 (5)	6 (7)	7 (3)	
Household finances (funds available at the end of the mo)				<.001
Has money left over	142 (47)	14 (17)	122 (60)	
Has just enough to make ends meet	80 (27)	29 (36)	47 (23)	
Does not have enough money to make ends meet	52 (17)	27 (33)	20 (10)	
Refused	26 (9)	11 (14)	13 (6)	
Educational level				<.001
Did not graduate from high school	24 (8)	14 (17)	8 (4)	
High school	85 (28)	32 (40)	50 (25)	
Some college or technical school	76 (25)	19 (23)	53 (26)	
College	55 (18)	8 (10)	43 (21)	
Graduate school	60 (20)	8 (10)	48 (24)	
ECOG performance score				.13
0	90 (30)	23 (28)	64 (32)	
1	32 (11)	6 (7)	23 (11)	
2	124 (41)	32 (40)	85 (42)	
3	46 (15)	17 (21)	26 (13)	
4	8 (3)	3 (4)	4 (2)	
Cancer type				.91
Breast	84 (28)	20 (25)	58 (29)	
Gastrointestinal	53 (18)	15 (19)	35 (17)	
Hematologic	37 (12)	4 (5)	31 (15)	
Lung	36 (12)	9 (11)	26 (13)	
Prostate	35 (12)	20 (25)	14 (7)	
Oropharynx	29 (10)	8 (10)	19 (9)	
Genitourinary	18 (6)	1 (1)	16 (8)	
Sarcoma	5 (2)	4 (5)	1 (0.5)	
Melanoma	2 (1)	0 (0)	2 (1)	
Glioblastoma	1 (0)	0 (0)	0 (0)	
Global distress index symptoms				
Lack of appetite	111 (37)	34 (42)	69 (34)	.22
Lack of energy	204 (68)	51 (63)	150 (69)	.30
Pain	164 (55)	50 (62)	101 (50)	.07
Drowsiness	171 (57)	47 (58)	113 (56)	.75
Constipation	94 (31)	30 (37)	55 (27)	.10
Dry mouth	137 (46)	43 (53)	83 (42)	.07
Sadness	116 (39)	29 (36)	80 (40)	.55
Worrying	170 (57)	47 (58)	115 (57)	.87
Irritability	121 (40)	32 (40)	80 (40)	.99
Nervousness	116 (39)	30 (37)	78 (39)	.81
Continue cancer treatment with probability of survival, %				.002
Nearly 100	140 (47)	48 (59)	86 (43)	
90-99	22 (7)	9 (11)	11 (5)	
50-89	34 (11)	3 (4)	28 (14)	
10-49	50 (17)	16 (20)	31 (15)	
1-9	9 (3)	0 (0)	9 (4)	
Almost 0	7 (2)	0 (0)	5 (2)	
No treatments	38 (13)	5 (6)	32 (16)	

ECOG indicates Eastern Cooperative Oncology Group.




FIGURE 3. Patient utilities for hospice services.

subscales, respectively; adjusted mean utilities: 2.31 vs 1.83; β coefficient, 0.51; 95% CI, 0.11-0.92 [$P = .01$]). This model did not include economic factors, which are considered separately below.

To describe the relation between perceived needs for hospice services and economic factors, which often are associated closely with race,⁶⁸ we used household finances as an indicator of economic status (not enough, enough, or more than enough money left at the end of the month). In unadjusted analysis, patients with fewer financial resources had greater perceived needs for hospice services (more than enough money left over, 1.51; enough money left over, 2.34; not enough money left over, 2.53). When the perceived needs of African-American and white patients were compared after adjusting for household finances, there was no significant difference (adjusted mean utilities: 2.38 vs 1.75; β coefficient, 0.24; 95% CI, 0.22-0.70 [$P = .31$]). Similarly, in a combined model that included household finances as well as all of the patient characteristics used in earlier models, African Americans did not have greater perceived needs for hospice services.

In subanalyses, we compared the perceived needs for hospice services of African-American patients ($n = 76$) and white patients ($n = 170$) whose preferences excluded them from hospice. In these groups, African-American patients had significantly greater perceived needs than white patients (total utilities: 2.34 vs 1.81; rank-sum test; $P = .006$). We also compared the perceived needs of African-American patients whose treatment preferences made them ineligible for hospice ($n = 76$) and white patients whose treatment preferences were consistent with hospice

($n = 32$). After controlling for health status by adjusting for age, ECOG-PS, dependencies in ADLs, and GDI physical and psychologic subscale scores, ineligible African-American patients had greater perceived needs for hospice services than eligible whites (adjusted means: 2.31 vs 1.73; β coefficient, 0.70; 95% CI, 0.05-1.35 [$P = .03$]). However, after adjusting for household finances, these 2 groups had similar utilities for hospice services (1.91 vs 1.83; β coefficient, -0.34 ; 95% CI, -0.56 -0.11 [$P = .27$]). Because of the small number of African-American patients whose preferences made them eligible for hospice ($n = 5$), a comparison of these patients' perceived needs with those of ineligible African Americans was not possible.

DISCUSSION

The Medicare Hospice Benefit and similar benefits of other insurers were designed to ensure that patients in the last 6 months of life have access to high-quality palliative care. However, African-American patients are less likely to use hospice than white patients. The results from the current study suggest that disparities in hospice use among patients with cancer may be the result of cancer treatment preferences among African-American patients that make them ineligible despite greater perceived needs for hospice care. In particular, this study offers 3 main insights into these disparities in hospice access.

First, we observed that African-American patients with cancer are more likely than white patients to want aggressive cancer treatment. This difference in preferences persists after adjusting for clinical and demographic characteristics. This result is important because, although other studies have observed racial differences in preferences for life-sustaining technologies,²⁶⁻²⁹ they have not examined preferences for cancer treatment. Preferences for aggressive cancer care, as well as other aggressive technologies, could be linked to medical distrust, quality of care, or communication issues between patients and providers.

Second, we observed that African-American patients with cancer had greater perceived needs for specific hospice services compared with white patients. This difference appears to be independent of clinical characteristics (eg, symptom burden, performance status) that are associated with greater needs. Furthermore, African-American

patients whose treatment preferences would have excluded them from hospice had greater perceived needs than white patients who would have been eligible.

Third, we observed that the greater perceived need for hospice services among African Americans was attributable largely to differences in self-reported finances. That is, the observed differences in perceived needs between African-American and white patients may reflect financial resources, because those who reported the fewest financial resources also reported the greatest need for services. This finding is consistent with other studies suggesting that the mechanisms driving health disparities include not only cultural differences associated with race but also economic characteristics that fundamentally may be unrelated to race.^{36,68,69} Thus, these economic factors should be explored further to determine why poorer patients want more services. For instance, it is possible that wealthier patients have more resources for care at the end of life and use these resources to obtain services outside of hospice. If so, then wealthier patients may be in a better position to avoid the 'terrible choice' requiring that they forgo treatment to receive hospice benefits. However, this is not known.

Together, these findings suggest that the hospice eligibility criteria of Medicare and other insurers requiring patients to give up cancer treatment contribute to racial disparities in hospice use. Moreover, these criteria do not select those patients with the greatest needs for hospice services. To the degree that this eligibility requirement prevents hospice use by the patients with the greatest needs, it fails to fulfill its purpose and should be reconsidered. It also is unfair: Other Medicare-supported services do not require that patients forgo 1 treatment to get another.⁷⁰ Instead, our study suggests that hospice access could be made fairer by using eligibility criteria that are more directly need-based. For instance, eligibility might be determined better by assessing needs for specific hospice services, such as pain or symptom management. This would make eligibility for hospice similar to eligibility for the Medicare Home Care Benefit, which perhaps is the closest Medicare program.⁷⁰

This study has several limitations. First, the results reported here are based on patients' self-reported needs for hospice services, but it is possible that these patients could not appreciate adequately how the services described could benefit them.⁶³ However, patients argu-

ably are the best judges of their own needs, even if those assessments are imperfect.

Second, this study assessed only 5 hospice services. It is possible that the finding of greater perceived needs reported here by African-American patients would not be true for other services that hospice provides (eg, delivery of medications or durable medical equipment). Although this is possible, there is no reason to believe that the pattern observed for the 5 services described here are unique.

Third, in this study, we did not assess actual choices regarding hospice enrollment, which may be influenced by factors such as the availability of family members who can assist with home care. A patient without adequate informal caregiving support and supervision at home may be less likely to enroll. Therefore, further research is needed to define which patient characteristics, such as the availability of family support, influence actual choices regarding hospice enrollment.

For 25 years, hospice services have been restricted to patients in the last 6 months of life who are willing to give up curative treatment.¹⁸ Although this restriction was intended to control costs, it has substantially reduced access to hospice for many patients who have needs for hospice services, and a disproportionate number of these patients are African American. This eligibility criterion should be reconsidered, and needs-based criteria should be considered to make hospice eligibility criteria both fairer and more consistent with eligibility criteria for other health insurance benefits.⁷⁰

Conflict of Interest Disclosures

Supported by American Cancer Society Mentored Scholar Research grant MRSMT-08-013-01-CPPB (to J.F.), the University of Pennsylvania's National Institutes of Health-funded Center of Excellence for Cancer Communication Research grant (to J.F.), grant R01CA109540 (to D.J.C.), a Veterans Administration Advanced Research Career Development Award, a Presidential Early Career Award for Scientists and Engineers (to D.J.C.), and by grants from the Abramson Cancer Center (to P.O.), the Center for Clinical Epidemiology and Biostatistics (to J.F.), and the Center for Health Equity Research and Promotion (to D.J.C. and J.F.).

References

1. Lutz S, Norrell R, Bertucio C, et al. Symptom frequency and severity in patients with metastatic or locally recurrent lung cancer: a prospective study using the Lung Cancer

- Symptom Scale in a community hospital. *J Palliat Med.* 2001;4:157-165.
2. Coyle N, Adelhardt J, Foley KM, Portenoy RK. Character of terminal illness in the advanced cancer patient: pain and other symptoms in the last 4 weeks of life. *J Pain Symptom Manage.* 1990;5:83-93.
 3. Vaino A, Auvinen A. Prevalence of symptoms among patients with advanced cancer: an international collaborative study. *J Pain Symptom Manage.* 1996;12:3-10.
 4. Hickok JT, Morrow GR, McDonald S, Bellg AJ. Frequency and correlates of fatigue in lung cancer patients receiving radiation therapy: implications for management. *J Pain Symptom Manage.* 1996;11:370-377.
 5. Twycross R, Harcourt J, Bergl S. A survey of pain in patients with advanced cancer. *J Pain Symptom Manage.* 1996;12:273-282.
 6. Harrington V, Lackey NR, Gates MF. Needs of caregivers of clinic and hospice cancer patients. *Cancer Nurs.* 1996;19:118-125.
 7. McMillan SC, Mahon M. The impact of hospice services on the quality of life of primary caregivers. *Oncol Nurs Forum.* 1994;21:1189-1195.
 8. Steele RG, Fitch MI. Needs of family caregivers of patients receiving home hospice care for cancer. *Oncol Nurs Forum.* 1996;23:823-828.
 9. Kane R, Wales J, Bernstein L, Leibowitz A, Kaplan S. A randomised controlled trial of hospice care. *Lancet.* 1984;1:890-894.
 10. Casarett D, Karlawish J, Crowley R, Mirsch T, Morales K, Asch DA. Improving use of hospice care in the nursing home: a randomized controlled trial. *JAMA.* 2005;294:211-217.
 11. Teno J, Clarridge B, Casey V, et al. Family perspectives on end-of-life care at the last place of care. *JAMA.* 2004;291:88-93.
 12. Miller S, Mor V, Wu N, Gozalo P, Lapane K. Does receipt of hospice care in nursing homes improve the management of pain at the end of life? *J Am Geriatr Soc.* 2002;50:507-515.
 13. Miller SC, Mor V, Teno J. Hospice enrollment and pain assessment and management in nursing homes. *J Pain Symptom Manage.* 2003;26:791-799.
 14. Seamark DA, Williams S, Hall M, Lawrence CJ, Gilbert J. Dying from cancer in community hospitals vs. hospice: closest lay carers' perceptions. *Br J Gen Pract.* 1998;48:1317-1321.
 15. Greiner K, Allen SP, Ahluwalia JS. Hospice usage by minorities in the last year of life: results from the National Mortality Followback Survey. *J Am Geriatr Soc.* 2003;51:970-978.
 16. Han B, Remsburg RE, Iwashyna TJ. Differences in hospice use between black and white patients during the period 1992 through 2000. *Med Care.* 2006;44:731-777.
 17. Johnson KS, Kuchibhatla M, Tanis D, Tulskey JA. Racial differences in hospice revocation to pursue aggressive care. *Arch Intern Med.* 2008;168:218-224.
 18. United States Congress. Medicare Hospice Regulations: 42 Code of Federal Regulations, Part 418. Washington, DC: Federal Register; 1996.
 19. Gibson R. Palliative care for the poor and disenfranchised: a view from the Robert Wood Johnson Foundation. *J R Soc Med.* 2001;94:486-489.
 20. United States Congress. Testimony Presented Before the Congress of the United States House of Representatives Committee on Government Reform. October 19, 1999. Available at: <http://www.dyingwell.org/uschrtest.htm>. Accessed May 25, 2007.
 21. Gelband H, Foley K. Improving Palliative Care for Cancer: Summary and Recommendations. Washington, DC: National Academies Press; 2001.
 22. Byock IR, Forman WB, Appleton M. Academy of hospice physicians' position statement on access to hospice and palliative care. *J Pain Symptom Manage.* 1996;11:69-70.
 23. McGorty EK, Bornstein BH. Barriers to physicians' decisions to discuss hospice: insights gained from the United States hospice model. *J Eval Clin Pract.* 2003;9: 363-372.
 24. Daugherty CK, Steensma DP. Overcoming obstacles to hospice care: an ethical examination of inertia and inaction. *J Clin Oncol.* 2002;20:2752-2755.
 25. Miller SC, Weitzen S, Kinzbrunner B. Factors associated with the high prevalence of short hospice stays. *J Palliat Med.* 2003;6:725-736.
 26. Blackhall L, Frank G, Murphy S, Michel V, Palmer J, Azen S. Ethnicity and attitudes towards life sustaining technology. *Soc Sci Med.* 1999;48:1779-1789.
 27. Caralis P, Davis B, Wright K, Marcial E. The influence of ethnicity and race on attitudes toward advance directives, life-prolonging treatments, and euthanasia. *J Clin Ethics.* 1993;4:155-165.
 28. Degenholtz H, Thomas SB, Miller MJ. Race and the intensive care unit: disparities and preferences for end-of-life care. *Crit Care Med.* 2003;31:S373-S378.
 29. Mitchell SL, Teno JM, Roy J, Kabumoto G, Mor V. Clinical and organizational factors associated with feeding tube use among nursing home residents with advanced cognitive impairment. *JAMA.* 2003;290:73-80.
 30. Neubauer B, Hamilton C. Racial differences in attitudes toward hospice care. *Hosp J.* 1990;6:37-48.
 31. Hwang S, Chang V, Cogswell J, Srinivas S, Kasimis B. Knowledge and attitudes toward end-of-life care in veterans with symptomatic metastatic cancer. *Palliat Support Care.* 2003;1:221-230.
 32. Taxis J. Attitudes, values, and questions of African Americans regarding participation in hospice programs. *J Hospice Palliat Nurs.* 2006;8:77-85.
 33. Cort M. Cultural mistrust and use of hospice care: challenges and remedies. *J Palliat Med.* 2004;7:63-71.
 34. Born W, Greiner K, Sylvia E, Butler J, Ahluwalia J. Knowledge, attitudes, and beliefs about end-of-life care among inner-city African Americans and Latinos. *J Palliat Med.* 2004;7:247-256.

35. Foley K, Gelband H. Improving Palliative Care for Cancer: Summary and Recommendations. Institute of Medicine and National Research Council. Washington, DC: National Academies Press; 2001.
36. Crawley L, Payne R, Bolden J, Payne T, Washington P, Williams S. Palliative and end-of-life care in the African American community. *JAMA*. 2000;284:2518-2521.
37. National Hospice and Palliative Care Organization. National Trend Summary 2005. Washington, DC: National Hospice and Palliative Care Organization; 2006.
38. Cornoni J, Ostfeld A, Taylor J, et al. Establishing populations for epidemiologic studies of the elderly: study design and methodology. *Aging*. 1993;5:27-37.
39. Portenoy RK, Thaler HT, Kornblith AB, et al. The Memorial Symptom Assessment Scale: an instrument for the evaluation of symptom prevalence, characteristics and distress. *Eur J Cancer*. 1994;30A:1326-1336.
40. Cella DF, Tulsky DS, Gray G. The Functional Assessment of Cancer Therapy scale: development and validation of the general measure. *J Clin Oncol*. 1993;11:570-579.
41. Sherbourne CD, Stewart AL. The MOS Social Support survey. *Soc Sci Med*. 1991;32:705-714.
42. Jenkinson C, Wright L, Coulter A. Criterion validity and reliability of the SF-36 in a population sample. *Qual Life Res*. 1994;3:7-12.
43. Conill C, Verger E, Salamero M. Performance status assessment in cancer patients. *Cancer*. 1990;65:1864-1866.
44. Sloan JA, Loprinzi CL, Laurine JA, et al. A simple stratification factor prognostic for survival in advanced cancer: the good/bad/uncertain index. *J Clin Oncol*. 2001;19:3539-3546.
45. Katz S, Akpom CA. Index of ADL. *Med Care*. 1976;14:116-118.
46. Mahoney FI, Barthel DW. Functional evaluation: the Barthel Index. *Md State Med J*. 1965;14:62-65.
47. Green P, Rao V. Conjoint measurement for quantifying judgmental data. *J Market Res*. 1971;8:355-363.
48. Green P. Hybrid models for conjoint analysis: an expository review. *J Market Res*. 1984;21:155-159.
49. Leigh T, MacKay D, Summers J. Reliability and validity of conjoint analysis and self-explicated weights: a comparison. *J Market Res*. 1984;21:456-462.
50. Moore W, Semenik R. Measuring preference with hybrid conjoint analysis: the impact of a different number of attributes in the master design. *J Business Res*. 1988;16:261-274.
51. Leung G, Chan S, Chau P, Chua S. Using conjoint analysis to assess patients' preferences when visiting emergency departments in Hong Kong. *Acad Emerg Med*. 2001;8:894-898.
52. Markham F, Diamond J, Hermansen C. The use of conjoint analysis to study patient satisfaction. *Eval Health Prof*. 1999;22:371-378.
53. Ryan M. A role for conjoint analysis in technology assessment in health care. *Int J Technol Assess Health Care*. 1999;15:443-457.
54. Ryan M, Farrar S. Using conjoint analysis to elicit preferences for health care. *BMJ*. 2000;320:1530-1533.
55. Shackley P, Slack R, Michaels J. Vascular patients' preferences for local treatment: an application of conjoint analysis. *J Health Serv Res Policy*. 2001;6:151-157.
56. Jan S, Mooney G, Ryan M, Bruggemann K, Alexander K. The use of conjoint analysis to elicit community preferences in public health research: a case study of hospital services in South Australia. *Aust N Z J Public Health*. 2000;24:64-70.
57. Shea JA, Asch DA, Johnson RF, et al. What predicts gastroenterologists' and surgeons' diagnosis and management of common bile duct stones? *Gastrointest Endosc*. 1997;46: 40-47.
58. Fraenkel L, Bogardus S, Wittink DR. Understanding patient preferences for the treatment of lupus nephritis with adaptive conjoint analysis. *Med Care*. 2001;39:1203-1216.
59. Srinivasan V, Jain A, Malhotra N. Improving predictive power of conjoint analysis by constrained parameter estimation. *J Market Res*. 1983;20:433-438.
60. Srinivasan V, Flaschbart P, Dajani J, Hartley R. Forecasting the effectiveness of work-trip gasoline conservation policies through conjoint analysis. *J Market Res*. 1981;45: 157-172.
61. Casarett D, Crowley R, Hirschman K. How should clinicians describe hospice to patients and their families? *J Am Geriatr Soc*. 2004;52:1923-1928.
62. Casarett D, Crowley R, Stevenson C, Xie S, Teno J. Making difficult decisions about hospice enrollment: what do patients and families want to know? *J Am Geriatr Soc*. 2005;53:249-254.
63. Rickerson E, Harrold J, Carroll J, Kapo J, Casarett D. Timing of hospice referral and families' perceptions of services: Are earlier hospice referrals better? *J Am Geriatr Soc*. 2005;53: 819-823.
64. Fried TR, Bradley EH, Towle VR, Allore H. Understanding the treatment preferences of seriously ill patients. *N Engl J Med*. 2002;346:1061-1066.
65. Casarett D, Van Ness P, O'Leary J, Fried T. Are patient preferences for life-sustaining treatment really a barrier to hospice enrollment for older adults with serious illness? *J Am Geriatr Soc*. 2006;54:472-478.
66. Sawtooth Software, Inc. ACA/Hierarchical Bayes Technical Paper. Sawtooth Software Technical Paper Series. Sequim, Wash: Sawtooth Software, Inc.; 2001.
67. Mickey J, Greenland S. A study of the impact of confounder-selection criteria on effect estimation. *Am J Epidemiol*. 1989;129:125-137.
68. Braveman P, Cubbin C, Egerter S, et al. Socioeconomic status in health research: one size does not fit all. *JAMA*. 2005;294:2879-2888.
69. Shavers V, Brown M. Racial and ethnic disparities in the receipt of cancer treatment. Review. *J Natl Cancer Inst*. 2002;94:334-357.
70. Centers for Medicare & Medicaid Services. Your Medicare Benefits. Available at: <http://www.medicare.gov/Publications/Pubs/pdf/10116.pdf> Accessed June 4, 2007.